

Tau contribution and precision measurement of θ_{23} at a neutrino factory

D. Indumathi and Nita Sinha¹

The Institute of Mathematical Sciences, Chennai 600 113, India.

Abstract. We discuss precision measurements of the leading atmospheric parameters at a standard neutrino factory. The oscillation of the muon and electron neutrinos (anti-neutrinos) to tau neutrinos (anti-neutrinos) adds to the muon events sample (both right sign and wrong sign) via leptonic decays of the taus produced through charge-current interactions in the detector. We focus on how this contribution affects a precision measurement of the atmospheric mixing parameters and the deviation of $\nu_\mu \leftrightarrow \nu_\tau$ mixing from maximality.

Keywords: neutrino factories, tau neutrinos, maximal mixing

PACS: 14.60.Pq, 14.60.Fg, 96.40.Tv

INTRODUCTION

A neutrino factory, with decays of μ^+ or μ^- beams in the straight sections of a storage ring, produces a spectrum of electron and muon neutrinos via $\mu^+ \rightarrow \bar{\nu}_\mu e^+ \nu_e$ and $\mu^- \rightarrow \nu_\mu e^- \bar{\nu}_e$. Charged-current (CC) interactions of the muon anti-neutrinos (neutrinos) in a detector lead to the production of μ^+ (μ^-) or the “right-sign” (RS) muons. The electron neutrinos (anti-neutrinos), on the other hand, can *oscillate* into muon neutrinos (anti-neutrinos) producing muons with charge opposite to that of the unoscillated case and get detected as “wrong sign” (WS) muons.

Most studies with neutrino factories[1] have focused on the use of WS events to pin down the unknown reactor angle θ_{13} , the CP violating phase δ_{CP} and the mass hierarchy. RS muons are useful for precision measurements of the atmospheric mixing angle θ_{23} and the mass squared difference Δm^2 , apart from understanding cross-section and flux uncertainties. In particular, they are sensitive probes of whether $\nu_\mu \leftrightarrow \nu_\tau$ mixing is maximal (i.e. $\theta_{23} = \pi/4$, referred to as maximal θ_{23}). Measurement of deviation from maximality is of significance in developing models for neutrino masses and mixings.

We address the little-studied issue of contamination of the (RS or WS) muon events sample from oscillations of the muon or electron neutrinos (anti-neutrinos) to tau neutrinos (anti-neutrinos), which, through CC interactions, result in tau leptons that decay to muons. We focus on how this contribution affects a precision measurement of θ_{23} and its deviation from maximal mixing.

The oscillation probability $P_{\mu\tau}$ is large even if θ_{13} vanishes, being driven by a nearly maximal θ_{23} . In spite of CC cross-section suppression for the massive tau produc-

tion, there is still a sizeable production rate and tau decay (with rate into muons of $\sim 17\%$) enhances the RS muon event rates, especially at small muon energies. Since the θ_{23} dependent terms in $P_{\mu\mu}$ and $P_{\mu\tau}$ come with opposite sign, the combination of muons from direct production and from tau decays marginally *decreases* the sensitivity of the event rates to this angle. Cuts imposed to remove the tau events drastically reduce the direct muon events as well, worsening the sensitivity to the oscillation parameters. Thus, the total muon events (direct muons and from tau decay) are less sensitive to the deviation from maximality. Neglect of the tau contribution will lead to an incorrect conclusion about the precision possible for the deviation from maximality.

THE INPUTS: FLUXES, KINEMATICS, CROSS-SECTIONS

The neutrino factory fluxes

We assume a basic muon storage ring configuration [2] with muon beam energy $E_b = 25$ GeV, with $n_\mu = 5 \times 10^{20}$ useful decays per year. We integrate the neutrino fluxes over the muon beam angle α , assuming a gaussian angular divergence of the muon beam around the z-axis with standard deviation [3] $\sigma = 0.1/\gamma$, where $\gamma = E_b/m_\mu$. This is then averaged over a small neutrino opening angle, $\theta' < \varepsilon = 0.3\sigma$ or roughly 0.1 mr. The resulting neutrino (or antineutrino) spectrum with gaussian angular spread averaged over ($\theta' < \varepsilon, 0 \leq \phi' \leq 2\pi$) at the baseline distance L from source to detector is given by,

$$\begin{aligned} \frac{dN_\mu}{dE_\nu} &\equiv \frac{1}{E_b} \frac{1}{\int d\Omega} \int d\Omega \left\langle \frac{dN_\mu}{dy d\Omega} \right\rangle_G \\ &= \frac{4n_\mu \gamma^4 y^2}{\pi L^2 E_b} \left\{ 3 - 4y\gamma^2 - \frac{\beta}{2} (3 - 8y\gamma^2)(1 + c_\varepsilon) e^{-\sigma^2/2} \right\} \end{aligned}$$

¹ presenting author

$$\begin{aligned}
& -\frac{1}{3}y(\gamma^2-1)\left[4+c_\varepsilon+c_\varepsilon^2+e^{-2\sigma^2}3c_\varepsilon(1+c_\varepsilon)\right], \\
\frac{dN_e}{dE_\nu} & \equiv \frac{1}{E_b} \frac{1}{\int d\Omega} \int d\Omega \left\langle \frac{dN_e}{dyd\Omega} \right\rangle_G \\
& = \frac{24n_\mu \gamma^4 y^2}{\pi L^2 E_b} \left\{ 1 - 2y\gamma^2 - \frac{\beta}{2} (1 - 4y\gamma^2) (1 + c_\varepsilon) e^{-\sigma^2/2} \right. \\
& \quad \left. - \frac{1}{6}y(\gamma^2-1)\left[4+c_\varepsilon+c_\varepsilon^2+e^{-2\sigma^2}3c_\varepsilon(1+c_\varepsilon)\right] \right\}, \quad (1)
\end{aligned}$$

where $y = E_\nu/E_b$, $c_\varepsilon, s_\varepsilon$ refer to $\cos \varepsilon, \sin \varepsilon$.

The kinematics

We focus on the spectrum of the final state muons and hence require the detailed kinematics of the CC interactions, in which either muons or taus are produced, with the latter decaying into muons, where again we use the differential decay rates (see Ref. [4] for details). In the laboratory frame, a neutrino of flavor $l = \mu$ or τ interacts with a nucleon and produces the corresponding charged lepton l at an angle θ_l w.r.t. the incident neutrino direction. In the case of ν_τ interactions, the tau is produced at a very forward angle while the azimuthal angle of the muon from tau decay ϕ_μ , is restricted by the decay kinematics. The available phase space is restricted in both direct muon and tau production due to the constraint on the available energy for the lepton: $E_- < E_l < E_+$; see Ref. [4] for the detailed expressions on the constraints. The effect of this pinching in available energy, for the case of a τ lepton being produced, can be seen in Fig. 1 where the final hadronic mass m_X is plotted as a function of E_τ . The notation is standard: $m_X^2 = W^2 = (p+q)^2$, where p, q are the nucleon and intermediate gauge boson 4-momenta in the laboratory frame. For a typical neutrino energy $E_\nu = 10$ GeV, allowed energy range for $\cos \theta_\tau = 0.91$ is tiny; tau leptons are hence produced in a very forward direction while the direct muons, due to their lighter mass, are less restricted.

The cross-sections

Since the energies of interest range from a few GeV to 25 GeV, the CC interactions include quasi-elastic (QE), resonance (Res) and deep inelastic (DIS) processes. We consider the double differential cross-sections,

$$\frac{d\sigma}{dE_l d\cos \theta_l} = \frac{G_F^2 \kappa^2}{2\pi} \frac{p_l}{M} \left\{ \sum_{i=1}^5 a_i W_i \right\}, \quad (2)$$

where G_F is the Fermi constant, $\kappa = M_W^2/(Q^2 + M_W^2)$ is the propagator factor with the W boson mass, M_W , p_l is the magnitude of three-momentum of the charged lepton produced and W_i are structure functions corresponding




FIGURE 1. Kinematics of ν_τ –nucleon CC interactions. The allowed parabolas of constant $\cos \theta_\tau$ in the m_X – E_τ plane are shown for $E_\nu = 10$ GeV. The ends of the parabolas (at $m_X = M$) give the limits of the tau energy.

to the general decomposition of the hadronic tensor. $W_{4,5}$ appear only for massive final leptons. We have,

$$\begin{aligned}
\sum_{i=1}^5 a_i W_i & = \left(2W_1 + \frac{m_l^2}{M^2} W_4 \right) (E_l - p_l \cos \theta_l) + W_2 (E_l + p_l \cos \theta_l) \\
& \quad \pm \frac{W_3}{M} (E_\nu E_l + p_l^2 - (E_\nu + E_l) p_l \cos \theta_l) - \frac{m_l^2}{M} W_5. \quad (3)
\end{aligned}$$

The detailed expressions for W_i are taken from Ref. [5] where the specific structure functions are listed for QE, Res and DIS leading order (LO) processes.

EVENT RATES IN A FAR-DETECTOR

Preliminaries

We assume that the neutrinos interact with a 50 kton iron detector such as the proposed INO/ICAL or MIND. Both μ^+ and μ^- beams with equal exposure are considered. While RS events are sensitive to deviations of θ_{23} from maximality, inclusion of WS events may only marginally worsen the results; however, the advantage in being “charge-blind” is significant, hence, all muon events are simply added. The generic number of muon events at a distance L , as a function of the observed muon energy E is,

$$\begin{aligned}
\mathcal{R}_{i,D}^b(E) & = K \int_{E_{\text{thr}}}^{E_{\text{max}}} dE_\nu \frac{dN_i(E_\nu, L)}{dE_\nu} \cdot P_{i\mu}(E_\nu, L) \int_0^1 d\cos \theta_\mu \\
& \quad \int_{E_-}^{E_+} dE_\mu \frac{d\sigma_\mu(E_\nu, E_\mu, \theta_\mu)}{dE_\mu d\cos \theta_\mu} \cdot R(E_\mu, E), \\
\mathcal{R}_{i,\tau}^b(E) & = K \int_{E_{\text{thr}}}^{E_{\text{max}}} dE_\nu \frac{dN_i(E_\nu, L)}{dE_\nu} \cdot P_{i\tau}(E_\nu, L) \int_{c_{\min}}^1 d\cos \theta_\tau \\
& \quad \int_{E_-}^{E_+} dE_\tau \frac{d\sigma_\tau(E_\nu, E_\tau, \theta_\tau)}{dE_\tau d\cos \theta_\tau} \int_{\text{restr.}} dE_\mu d\cos \theta_\mu d\phi_\mu \\
& \quad \frac{1}{\Gamma_\tau} \frac{d\Gamma(E_\tau, E_\mu, \theta_\tau, \theta_\mu, \phi_\mu)}{dE_\mu d\cos \theta_\mu d\phi_\mu} \cdot R(E_\mu, E), \quad (4)
\end{aligned}$$




FIGURE 2. Muon event rates as a function of the observed muon energy. RS and WS events from μ^- and μ^+ beams are shown in the upper panels. Contributions from direct muon production (denoted by D) and that of muons from tau decay (labeled as τ) are separately shown. The left lower panel shows the sum $D + \tau$. Oscillation parameters are as given in text.

where the superscript b refers to the beam-type, $i = \mu, \bar{e}(\bar{\mu}, e)$ denotes the neutrino type from $\mu^- (\mu^+)$ beam. Here $K = N_t n_y$, N_t is number of target nucleons (assumed isoscalar), n_y is number of years of data and R is the Gaussian energy resolution function of width $\sigma = 0.07 E_\mu$. The subscripts D and τ correspond to the production of direct muons and τ production and decay into muons, respectively. The limits of integration and the restriction from an angular constraint, in case of tau decay are defined in Ref. [4]. The charge-blind events are obtained by summing over the index i and the total events for each beam are then obtained by the sum, $\mathcal{R}_{\text{tot}}^b(E) = \sum_i (\mathcal{R}_{i,D}^b + \mathcal{R}_{i,\tau}^b(E))$. We assume 90% reconstruction efficiency of muons. Muon event rates accumulated over five years are used to study the sensitivity to the deviation of the mixing angle θ_{23} from maximality.

Typical event rates at $L=7400$ Km (magic baseline) for oscillation parameters, $\Delta m^2 = 2.4 \times 10^{-3} \text{ eV}^2$, $\theta_{23} = 42^\circ$, $\theta_{13} = 1^\circ$, $\sin^2 \theta_{12} = 0.304$ and $\Delta_{21} = 7.65 \times 10^{-5} \text{ eV}^2$, are shown as a function of the observed lepton energy in Fig. 2. The panels show the direct muon production and tau decay contributions to the RS and WS muon events from μ^- and μ^+ beams. Since the neutrino-nucleon cross-sections are larger than those for anti-neutrinos, the $\mu^- (\mu^+)$ beams are preferred for studying the RS (WS) events where the cross-sections are larger. However, the full sample without charge identification is preferable, due to higher detection efficiency. We therefore add the events (RS+WS) from both μ^- and μ^+ beams.

It can be seen that there is a substantial contribution




FIGURE 3. Effects of angular cuts on the tau contribution to muon events at neutrino factories. For details, see the text.

to the RS events from tau decay into muons. The tau contribution alters the sensitivity to the oscillation parameters; we next discuss whether these events can be removed through suitable cuts.

Cuts on tau contribution

Since tau production in neutrino-nucleon interactions is extremely forward-peaked, one obvious way to remove the tau contribution is with an angular cut. Also, since the tau contribution is substantial at small observed muon energies where the tau decay rate is large, a muon energy cut can also be contemplated. The effect of cuts on the event rates is seen in Fig. 3 – the only cut effective in removing the tau contribution is one ($\theta_\mu > 25^\circ$) that removes the signal itself! Alternately a muon energy cut of $E > 10 - 15 \text{ GeV}$ can substantially remove the tau contribution, still leaving sufficient direct muons. However, such a large energy cut will worsen the measured precision of the mixing parameters as sensitivity is higher in the lower energy bins where matter effects are large. In short, it is not feasible to cut out the tau contribution and still make a precision measurement, in this case, of the deviation of θ_{23} from maximality.

Effect of the tau contribution

As stated earlier, the tau contribution alters the dependence on the mixing parameters, altering the precision to which we can determine them. While the tau events have less sensitivity to θ_{23} , the rate increases while the direct event rate decreases, as θ_{23} increases towards max-

imal $\theta_{23} = \pi/4$. The inclusion of muons from tau events also alters the uncertainties considerably. A near detector sensitive to muons, measures the combination of flux times cross-section of the muons. This also appears in the RS event rate for direct muon production and is therefore well constrained. However, for tau production and decay, the RS event rate has the combination of muon flux and the tau production cross-section. The heavy tau cross-sections have larger uncertainties, where mass corrections are large. Furthermore, since these contributions result from oscillations, no near detector can help reduce the uncertainties. Hence overall uncertainties are much larger for the tau contribution than for direct muons.

Hence, in our numerical calculations we use an overall normalization error of 0.1% for direct, while a modest 2% is used for the total (direct+tau), muon events. We use typical input values of $(\Delta m^2, \theta_{23}, \theta_{13})$ to estimate how well the generated “data” can be fitted, and calculate the resulting precision on the parameters. We keep the solar parameters fixed at their best-fit values of Ref. [6] and set δ_{CP} to zero. The best fits (and regions of confidence levels in parameter space) are obtained by minimizing the chi-squared with a pull corresponding to the normalization uncertainties specified.

We present results for a typical sample set of input parameters, $(\Delta m^2, \theta_{23}, \theta_{13} = 2.4 \times 10^{-3} \text{ eV}^2, 41.9^\circ, 1^\circ)$. We minimize first over the pull, and then over Δm^2 and θ_{23} , keeping θ_{13} fixed. Fig. 4 shows the allowed Δm^2 – θ_{23} parameter space at 99% CL. The solid and dashed lines correspond to considering direct and total (including those from tau decay) muon events respectively. Note that the 99% CL contour is much more constrained with direct than for total muons. In particular, it is the Δm^2 values that are smaller than the input value, that broaden the contour and limit the discrimination. The largest true value of θ_{23} that can be discriminated from maximal is shown in Fig. 5, as a function of Δm^2 again, for $\theta_{13} = 1^\circ$. It is seen that tau contamination worsens the ability to discriminate θ_{23} from maximal, thus making this measurement harder than originally expected.

CONCLUSION

The oscillations of the muon or electron neutrinos (anti-neutrinos) to tau neutrinos (anti-neutrinos) results in tau leptons produced through CC interactions in the detector which on leptonic decay add to the right as well as wrong sign muon events obtained directly. This tau contamination worsens the ability to discriminate against maximal $\nu_\mu \leftrightarrow \nu_\tau$ mixing. It is practically impossible to devise satisfactory cuts to remove this tau contamination. Uncertainties from this tau background must be brought under control before making precision parameter measurements at neutrino factories.




FIGURE 4. Allowed Δm^2 – θ_{23} parameter space at 99% CL from CC muon events, directly produced (solid line) and with inclusion of those from tau decay (dashed line). See text for values of input parameters.




FIGURE 5. The largest (smallest) true value of θ_{23} in the first (second) octant that can be discriminated from $\theta_{23} = \pi/4$, as a function of Δm^2 are shown when only Direct (D) and total (D + τ) events are considered; Here θ_{13} is fixed at 1° .

ACKNOWLEDGMENTS

NS thanks the organizers for a wonderful conference as well as for local hospitality.

REFERENCES

1. A. Bandyopadhyay *et al.* [ISS Physics Working Group], Rept. Prog. Phys. **72**, 106201 (2009) and references therein.
2. IDS-NF baseline study, IDS-NF-Baseline-2007/1.0, prepared by the IDS-NF Steering Group, Nov 2008.
3. C. Crisan and S. Geer, Fermilab-TM-2101; A. Broncano and O. Mena, Eur. Phys. J. C **29**, 197 (2003) and Ref. [1].
4. D. Indumathi and N. Sinha, arXiv:0910.2020 [hep-ph].
5. K. Hagiwara, K. Mawatari and H. Yokoya, Nucl. Phys. B **668**, 364 (2003) [Erratum-ibid. B **701**, 405 (2004)].
6. T. Schwetz *et al.*, New J. Phys. **10** 113011 (2008).